

Ground-State Energy and Condensate Density of a Dilute Bose Gas Revisited

Kazumasa TSUTSUI and Takafumi KITA

Department of Physics, Hokkaido University, Sapporo 060-0810, Japan

(Dated: May 6, 2013)

The ground-state energy per particle E/N and condensate density n_0 of a dilute Bose gas are studied with a self-consistent perturbation expansion satisfying the Hugenholtz-Pines theorem and conservation laws simultaneously. A class of Feynman diagrams for the self-energy, which has escaped consideration so far, is shown to add an extra constant $c_{ip} \sim O(1)$ to the expressions reported by Lee, Huang, and Yang [Phys. Rev. **106** (1957) 1135] as $E/N = (2\pi\hbar^2 an/m)[1 + (128/15\sqrt{\pi} + 16c_{ip}/5)\sqrt{a^3n}]$ and $n_0/n = 1 - (8/3\sqrt{\pi} + c_{ip})\sqrt{a^3n}$, where a , n , and m are the s -wave scattering length, particle density, and particle mass, respectively. We present a couple of estimates for c_{ip} ; the third-order perturbation expansion yields $c_{ip} = 0.412$.

Lee *et al.*¹ studied the ground-state properties of a single-component homogeneous dilute Bose gas based essentially on Bogoliubov theory² at zero temperature. They obtained widely accepted expressions for the energy per particle and condensate density as^{1,3,4}

$$\frac{E}{N} = \frac{2\pi\hbar^2 an}{m} \left(1 + \frac{128}{15\sqrt{\pi}} \sqrt{a^3n} \right), \quad (1a)$$

$$\frac{n_0}{n} = 1 - \frac{8}{3\sqrt{\pi}} \sqrt{a^3n}, \quad (1b)$$

where a , n , and m are the s -wave scattering length, particle density, and particle mass, respectively. On the other hand, a standard finite-temperature extension of Bogoliubov theory is known to yield an unphysical energy gap in the single-particle spectrum^{5,6} in contradiction to the Hugenholtz-Pines theorem.⁷ This issue was discussed extensively by Hohenberg and Martin⁸ in connection with conservation laws.

Recently, a self-consistent perturbation expansion has been formulated for condensed Bose systems so that the Hugenholtz-Pines theorem and conservation laws are satisfied simultaneously.^{9,10} A notable prediction of this theory¹¹ is that there is a new class of Feynman diagrams for the self-energy, i.e., those that may be classified as *improper* or *reducible* following the conventional terminology and should certainly be excluded from its definition in the normal state.^{3,4,12} The rationale for their existence in condensed Bose systems is that they are indispensable for three exact statements, i.e., the Hugenholtz-Pines theorem, conservation laws, and an identity for the interaction energy, to be satisfied order by order simultaneously in the self-consistent perturbation expansion.⁹ Thus, the conventional procedure of constructing the self-energy in the Dyson-Beliaev equation with the concept of *properness* (or *irreducibility*)^{3,4} cannot be justified. Since they have been overlooked, it will be worth looking at how the *improper* diagrams affect the standard results for the dilute Bose gas.^{3,4}

In the present paper, we will show that the *improper* diagrams change eq. (1) into

$$\frac{E}{N} = \frac{2\pi\hbar^2 an}{m} \left[1 + \frac{16}{5} \left(\frac{8}{3\sqrt{\pi}} + c_{ip} \right) \sqrt{a^3n} \right], \quad (2a)$$

$$\frac{n_0}{n} = 1 - \left(\frac{8}{3\sqrt{\pi}} + c_{ip} \right) \sqrt{a^3n}, \quad (2b)$$

where c_{ip} is a numerical constant of order 1. We will subsequently estimate c_{ip} using a couple of approximations. One of them, i.e., the perturbation expansion up to the third order, will be shown to yield $c_{ip} = 0.412$.

We consider a homogeneous system of identical Bose particles with mass m and spin 0 interacting via the contact potential $U\delta(\mathbf{r}_1 - \mathbf{r}_2)$. The Hamiltonian is given by

$$H = \sum_{\mathbf{p}} (\epsilon_{\mathbf{p}} - \mu) c_{\mathbf{p}}^\dagger c_{\mathbf{p}} + \frac{U}{2V} \sum_{\mathbf{p}_1 \mathbf{p}_2 \mathbf{q}} c_{\mathbf{p}_1 + \mathbf{q}}^\dagger c_{\mathbf{p}_2 - \mathbf{q}}^\dagger c_{\mathbf{p}_2} c_{\mathbf{p}_1}, \quad (3)$$

where $\mathbf{p}$, $\epsilon_{\mathbf{p}}$, μ , and V are the momentum, kinetic energy, chemical potential, and volume, and $c_{\mathbf{p}}^\dagger$ and $c_{\mathbf{p}}$ are the creation and annihilation operators, respectively. It is convenient to set $\hbar = 2m = k_B = T_c^0 = 1$, where k_B denotes the Boltzmann constant and T_c^0 is the transition temperature of ideal Bose-Einstein condensation.⁴ Thus, the kinetic energy is expressed simply as $\epsilon_{\mathbf{p}} = p^2$. Ultra-violet divergences inherent in the continuum model are removed here by introducing a momentum cutoff $p_c \gg 1$. However, our final results will be free from p_c , as seen below. It is standard in the low-density limit to remove U in favor of the s -wave scattering length a . They are connected in the conventional units by

$$\frac{m}{4\pi\hbar^2 a} = \frac{1}{U} + \int \frac{d^3p}{(2\pi\hbar)^3} \frac{\theta(p_c - p)}{2\epsilon_p},$$

with $\theta(x)$ the step function, which in the present units reads $1/8\pi a = 1/U + p_c/4\pi^2$. We will focus on the limit $a \rightarrow 0$ and choose p_c so that $1 \ll p_c \ll \pi/2a$ is satisfied. Thus, we can set

$$U = \frac{8\pi a}{1 - (2/\pi)p_c a} \approx 8\pi a \left(1 + \frac{2}{\pi} p_c a \right) \quad (4)$$

as an excellent approximation for the present purpose.

Let us recapitulate some relevant results from ref. 9. Green's function for the condensed Bose system can be expressed in the Nambu representation as

$$\hat{G}_{\vec{p}} \equiv \begin{bmatrix} G_{\vec{p}} & F_{\vec{p}} \\ -\bar{F}_{\vec{p}} & -\bar{G}_{\vec{p}} \end{bmatrix}, \quad (5a)$$

where $\vec{p} = (\mathbf{p}, i\varepsilon_\ell)$ with $\varepsilon_\ell \equiv 2\ell\pi T$ ($\ell = 0, \pm 1, \pm 2, \dots$) the Matsubara frequency. The upper elements satisfy $G_{\vec{p}} = G_{\vec{p}}^*$ and $F_{\vec{p}} = F_{-\vec{p}}$, and a barred quantity generally denotes $\bar{G}_{\vec{p}} \equiv G_{-\vec{p}}^*$. The matrix Green's function obeys the Dyson-Beliaev equation

$$\hat{G}_{\vec{p}} = \begin{bmatrix} i\varepsilon_\ell - \epsilon_p - \Sigma_{\vec{p}} + \mu & -\Delta_{\vec{p}} \\ \bar{\Delta}_{\vec{p}} & i\varepsilon_\ell + \epsilon_p + \bar{\Sigma}_{\vec{p}} - \mu \end{bmatrix}^{-1}, \quad (5b)$$

which may also be regarded as defining the self-energies $\Sigma_{\vec{p}}$ and $\Delta_{\vec{p}}$. In the self-consistent perturbation expansion, they are obtained from a functional $\Phi[\hat{G}_{\vec{p}}, n_0]$ as

$$\Sigma_{\vec{p}} = -T^{-1} \frac{\delta \Phi}{\delta G_{\vec{p}}}, \quad \Delta_{\vec{p}} = 2T^{-1} \frac{\delta \Phi}{\delta \bar{F}_{\vec{p}}}. \quad (6a)$$

In addition, Φ satisfies

$$\frac{1}{V} \frac{\delta \Phi}{\delta n_0} = \Sigma_{\vec{0}} - \Delta_{\vec{0}}. \quad (6b)$$

The substitution of eq. (6a) into eq. (5b) yields self-consistent (i.e., nonlinear) equations for $G_{\vec{p}}$ and $F_{\vec{p}}$. It also follows from eq. (6b) that the stationarity condition $\delta\Omega/\delta n_0 = 0$ for the thermodynamic potential Ω is equivalent to the Hugenholtz-Pines relation

$$\mu = \Sigma_{\vec{0}} - \Delta_{\vec{0}}. \quad (7)$$

These are exact statements. It has been shown that the key functional Φ can be constructed as a power-series expansion in U in such a way that eq. (7), conservation laws, and an exact relation for the interaction energy are fulfilled simultaneously order by order.

The first-order functional $\Phi^{(1)}$ is given graphically in Fig. 1, which can be shown to reproduce eq. (1) as follows. The differentiations of eq. (6a) correspond to removing a line of $G_{\vec{p}}$ and $\bar{F}_{\vec{p}}$, respectively, from every diagram in Fig. 1 in all possible ways. Hence, $\Sigma^{(1)}$ and $\Delta^{(1)}$ are obtained as

$$\Sigma^{(1)} = 2Un, \quad (8a)$$

$$\Delta^{(1)} = U \left(n_0 - \sum_{\vec{p}} F_{\vec{p}}^{(1)} \right), \quad (8b)$$




FIG. 1: Feynman diagrams for $\Phi^{(1)}$. A filled circle denotes $2U$, a line with an arrow (two arrows) represents G (either F or $\bar{F}$) in eq. (5a) as in the theory of superconductivity,^{3,4} and every missing line in the last three diagrams corresponds to n_0 . The number below each diagram indicates its relative weight, which should be multiplied by $1/4$ to obtain the absolute weight.

where

$$n = n_0 - \sum_{\vec{p}} G_{\vec{p}}^{(1)} e^{i\varepsilon_\ell 0+} \quad (8c)$$

is the particle density with 0_+ an infinitesimal positive constant,^{3,12} and the summation over $\vec{p}$ denotes

$$\sum_{\vec{p}} \equiv T \sum_{\ell=-\infty}^{\infty} \int \frac{d^3p}{(2\pi)^3} \xrightarrow{T \rightarrow 0} \int_{-\infty}^{\infty} \frac{d\varepsilon_\ell}{2\pi} \int \frac{d^3p}{(2\pi)^3}.$$

Thus, the first-order self-energies $\Sigma^{(1)}$ and $\Delta^{(1)}$ have no $\vec{p}$ dependence, and eq. (7) reduces to

$$\mu^{(1)} = \Sigma^{(1)} - \Delta^{(1)}. \quad (8d)$$

Substituting eqs. (8a), (8b), and (8d) into eq. (5b) and carrying out matrix inversion, we obtain the upper elements of $\hat{G}_{\vec{p}}$ in eq. (5a) as

$$\begin{bmatrix} G_{\vec{p}}^{(1)} \\ F_{\vec{p}}^{(1)} \end{bmatrix} = \frac{-1}{\varepsilon_\ell^2 + \epsilon_p(\epsilon_p + 2\Delta^{(1)})} \begin{bmatrix} i\varepsilon_\ell + \epsilon_p + \Delta^{(1)} \\ \Delta^{(1)} \end{bmatrix}. \quad (9)$$

With eq. (9), we can perform the summations over $\vec{p}$ in eqs. (8b) and (8c) analytically to obtain

$$\Delta^{(1)} = U \left[n_0 + \frac{\Delta^{(1)}}{4\pi^2} p_c - \frac{(2\Delta^{(1)})^{3/2}}{8\pi^2} \right], \quad (10a)$$

$$n = n_0 + \frac{(2\Delta^{(1)})^{3/2}}{24\pi^2}, \quad (10b)$$

where we have set $p_c \rightarrow \infty$ in the second convergent integral; this procedure will be adopted throughout. The substitution of eq. (4) into eq. (8a) yields

$$\Sigma^{(1)} = 16\pi an \left(1 + \frac{2}{\pi} p_c a \right). \quad (11a)$$

For $\Delta^{(1)}$, eqs. (4) and (10) indicate that $\Delta^{(1)} \approx 8\pi an$ to the leading order. Collecting next-order terms perturbatively with $p_c a \ll 1$ in mind, we arrive at

$$\Delta^{(1)} = 8\pi an \left(1 + \frac{4}{\pi} p_c a - \frac{32}{3\sqrt{\pi}} \sqrt{a^3 n} \right). \quad (11b)$$

Let us substitute eq. (11) into eqs. (8d) and (10b). We then obtain

$$\mu^{(1)} = 8\pi an \left(1 + \frac{32}{3\sqrt{\pi}} \sqrt{a^3 n} \right) \quad (12)$$

and eq. (1b), respectively, which are correct up to the next-to-the-leading order and also free from p_c . Finally integrating the thermodynamic relation $\mu = \partial E / \partial N$ over the particle number N , we arrive at eq. (1a) in units of $\hbar = 2m = 1$. Thus, the Lee-Huang-Yang results of eq. (1) have been reproduced in our mean-field approximation.




FIG. 2: Feynman diagrams for $\Phi^{(2)}$. A filled circle denotes $2U$, a line with an arrow (two arrows) represents G (either F or $\bar{F}$) in eq. (5a), and every missing line in the second and third rows corresponds to n_0 . The number below each diagram indicates its relative weight, which should be multiplied by $-1/8$ to obtain the absolute weight.

Now, we proceed to look into higher-order terms. The second-order $\Phi^{(2)}$ can also be constructed uniquely so as to satisfy eq. (7), conservation laws, and an exact relation for the interaction energy.⁹ The results are shown graphically in Fig. 2. For the dilute Bose gas at $T = 0$, in general, the relative importance of each diagram increases as the number of Green's function lines in it increases, as our previous analysis on eq. (8) indicates. Hence, the dominant contribution of the second order originates from the third-row diagrams in Fig. 2, which also bring a novel structure to the self-energies. Indeed, the differentiations of eq. (6a) for these diagrams yield

$$\Sigma_{\vec{p}}^{(2ip)} = \Delta_{\vec{p}}^{(2ip)} = 2(U n_0)^2 (G_{\vec{p}} + \bar{G}_{\vec{p}} - F_{\vec{p}} - \bar{F}_{\vec{p}}), \quad (13a)$$

which is classified as *improper* or *reducible* in the normal state^{3,4,12} and hence has been overlooked so far. This structure contains nothing that contradicts eq. (5b), which defines self-energies, however. It is a natural consequence of the requirement that the Hugenholtz-Pines relation (7), which is an exact statement, is obeyed order by order in the self-consistent perturbation expansion. We will show shortly that it brings an additional term to eq. (1). On the other hand, the diagrams in the first and second rows in Fig. 2, which yield conventional *proper* or *irreducible* self-energies, are at least one order of magnitude smaller at $T = 0$; thus, they can be neglected for the present purpose.

The *improper* structure extends beyond the second order. In the third order, for example, the rightmost diagram in Fig. 3 also yields^{9,11}

$$\Sigma_{\vec{p}}^{(3ip)} = \Delta_{\vec{p}}^{(3ip)} = \frac{5}{4} (2U n_0)^3 (G_{\vec{p}} + \bar{G}_{\vec{p}} - F_{\vec{p}} - \bar{F}_{\vec{p}})^2, \quad (13b)$$

where the factor $5/4$ is the sum of 1 and $1/2^2$ originating from the particle-hole and particle-particle bubble diagrams, respectively. Equation (13) reveals common features of the leading-order *improper* contribution at each order: (i) they are functionals of

$$f_{\vec{p}} \equiv G_{\vec{p}} + \bar{G}_{\vec{p}} - F_{\vec{p}} - \bar{F}_{\vec{p}}, \quad (14)$$



FIG. 3: Feynman diagrams for $\Phi^{(3)}$ drawn without arrows.

and (ii) the diagonal and off-diagonal self-energies are the same. Note the symmetry $f_{\vec{p}} = \bar{f}_{\vec{p}}$.

Adding the series of eq. (13) to eqs. (8a) and (8b), we can express the diagonal and off-diagonal self-energies as

$$\Sigma_{\vec{p}} = \Sigma^{(1)} + \Delta_{\vec{p}}^{(ip)}, \quad \Delta_{\vec{p}} = \Delta^{(1)} + \Delta_{\vec{p}}^{(ip)}, \quad (15)$$

respectively, where $\Delta_{\vec{p}}^{(ip)} \equiv \Delta_{\vec{p}}^{(2ip)} + \Delta_{\vec{p}}^{(3ip)} + \dots$ satisfies $\Delta_{\vec{p}}^{(ip)} = \bar{\Delta}_{\vec{p}}^{(ip)}$. The corresponding Green's functions are obtained by using eqs. (5) and (7) as

$$\begin{bmatrix} G_{\vec{p}} \\ F_{\vec{p}} \end{bmatrix} = \frac{-1}{\varepsilon_{\vec{p}}^2 + \epsilon_p(\epsilon_p + 2\Delta_{\vec{p}})} \begin{bmatrix} i\varepsilon_{\vec{p}} + \epsilon_p + \Delta_{\vec{p}} \\ \Delta_{\vec{p}} \end{bmatrix}, \quad (16)$$

with

$$\mu = \Sigma^{(1)} - \Delta^{(1)}. \quad (17a)$$

This expression for μ appears to be the same as eq. (8d), but the right-hand side now should be calculated by using eq. (16) instead of eq. (9) in the self-consistent approximation scheme. To be specific, $\Sigma^{(1)}$ is still given by eq. (8a) but with $G_{\vec{p}}^{(1)} \rightarrow G_{\vec{p}}$ in eq. (8c), i.e.,

$$n = n_0 - \sum_{\vec{p}} G_{\vec{p}}^{(1)} e^{i\varepsilon_{\vec{p}} 0^+} - \sum_{\vec{p}} (G_{\vec{p}} - G_{\vec{p}}^{(1)}). \quad (17b)$$

This redundant expression enables us to treat the third term on the right-hand side as a correction to the mean-field result (1b). Similarly, eq. (8b) is replaced by

$$\Delta^{(1)} = U \left[n_0 - \sum_{\vec{p}} F_{\vec{p}}^{(1)} - \sum_{\vec{p}} (F_{\vec{p}} - F_{\vec{p}}^{(1)}) \right]. \quad (17c)$$

On the other hand, the substitution of eq. (16) into the series of eq. (13) yields the equation for $\Delta_{\vec{p}}^{(ip)} \equiv \Delta_{\vec{p}}^{(2ip)} + \Delta_{\vec{p}}^{(3ip)} + \dots$ as

$$\Delta_{\vec{p}}^{(ip)} = U n_0 \left[-\frac{2U n_0}{\eta_{\vec{p}} + \Delta_{\vec{p}}^{(ip)}} + \frac{10(U n_0)^2}{(\eta_{\vec{p}} + \Delta_{\vec{p}}^{(ip)})^2} + \dots \right], \quad (18)$$

with

$$\eta_{\vec{p}} \equiv \frac{\varepsilon_{\vec{p}}^2 + \epsilon_p(\epsilon_p + 2\Delta^{(1)})}{2\epsilon_p}. \quad (19)$$

Equation (18) forms an algebraic equation for $\Delta_{\vec{p}}^{(ip)}$, and we see immediately that $\Delta_{\vec{p}}^{(ip)} \sim U n_0 \sim U n$, i.e., it is of the same order as $\Delta^{(1)}$ in eq. (8b).

Let us transform the third term on the right-hand side of eq. (17b) by using eqs. (9) and (16), noting the summation over $\vec{p}$ below eq. (8c) at $T = 0$, and making a change of variables as

$$\epsilon_p = Un\tilde{\epsilon}_p, \quad \varepsilon_\ell = Un\tilde{\varepsilon}_\ell, \quad \Delta_{\vec{p}} = Un\tilde{\Delta}_{\vec{p}}. \quad (20)$$

The result may be written as

$$\sum_{\vec{p}} \left(G_{\vec{p}} - G_{\vec{p}}^{(1)} \right) = - \left(\frac{Un}{8\pi} \right)^{3/2} c_{\text{ip}}, \quad (21)$$

where c_{ip} is given by

$$c_{\text{ip}} \equiv \frac{4\sqrt{2}}{\pi^{3/2}} \int_0^\infty d\tilde{\epsilon}_p \tilde{\epsilon}_p^{1/2} \int_0^\infty d\tilde{\varepsilon}_\ell \frac{(\tilde{\varepsilon}_\ell^2 - \tilde{\epsilon}_p^2) \tilde{\Delta}_{\vec{p}}^{(\text{ip})}}{[\tilde{\varepsilon}_\ell^2 + \tilde{\epsilon}_p(\tilde{\epsilon}_p + 2\tilde{\Delta}_{\vec{p}})][\tilde{\varepsilon}_\ell^2 + \tilde{\epsilon}_p(\tilde{\epsilon}_p + 2\tilde{\Delta}_{\vec{p}}^{(1)})]}. \quad (22)$$

When substituting eq. (21) into eq. (17b) for the present purpose, we can replace $(Un/8\pi)^{3/2}$ by the leading-order expression $(na)^{3/2}$ with $\tilde{\Delta}_{\vec{p}}^{(1)} \rightarrow 1$ in eq. (22); see eqs. (4), (8), and (10) on this point. Equation (18) is also approximated with $Un_0 \approx Un$ as

$$\tilde{\Delta}_{\vec{p}}^{(\text{ip})} = -\frac{2}{\tilde{\Delta}_{\vec{p}}^{(\text{ip})} + \tilde{\eta}_{\vec{p}}} + \frac{10}{(\tilde{\Delta}_{\vec{p}}^{(\text{ip})} + \tilde{\eta}_{\vec{p}})^2} + \dots, \quad (23)$$

where $\tilde{\eta}_{\vec{p}}$ is defined by

$$\tilde{\eta}_{\vec{p}} \equiv (\tilde{\varepsilon}_\ell^2 + \tilde{\epsilon}_p^2)/2\tilde{\epsilon}_p + 1. \quad (24)$$

Equation (23) with eq. (24) implies that $\tilde{\Delta}_{\vec{p}}^{(\text{ip})}$ is a function of only $x \equiv \tilde{\varepsilon}_\ell^2/\tilde{\epsilon}_p + \tilde{\epsilon}_p$. Hence, it is convenient to make a further change of variables in eq. (22):

$$x \equiv \frac{\tilde{\varepsilon}_\ell^2}{\tilde{\epsilon}_p} + \tilde{\epsilon}_p, \quad \xi \equiv \tilde{\epsilon}_p, \quad b(x) \equiv \frac{x}{2} + 1, \quad y(x) \equiv \tilde{\Delta}_{\vec{p}}^{(\text{ip})}, \quad (25)$$

to transform it to

$$c_{\text{ip}} = \frac{2\sqrt{2}}{\pi^{3/2}} \int_0^\infty d\xi \int_\xi^\infty dx \frac{(x - 2\xi)y(x)}{\sqrt{x - \xi}[x + 2 + 2y(x)](x + 2)} \\ = -\frac{4\sqrt{2}}{3\pi^{3/2}} \int_0^\infty dx \frac{x^{3/2}y(x)}{[x + 2 + 2y(x)](x + 2)}. \quad (26)$$

The last expression has been obtained by changing the order of integrations and subsequently integrating over ξ . A similar analysis on the third term in the square brackets of eq. (17c) yields

$$\sum_{\vec{p}} \left(F_{\vec{p}} - F_{\vec{p}}^{(1)} \right) = \left(\frac{Un}{8\pi} \right)^{3/2} c_F, \quad (27)$$

where c_F is the same as eq. (22) except for $\tilde{\varepsilon}_\ell^2 - \tilde{\epsilon}_p^2 \rightarrow -\tilde{\varepsilon}_\ell^2 - \tilde{\epsilon}_p^2$ in the numerator. Transforming it in the same manner as from eq. (22) into eq. (26), we find that

$$c_F = 3c_{\text{ip}}. \quad (28)$$

Let us substitute eqs. (21) and (27) with eq. (28) into eq. (17). We then find that the *improper* contribution changes eqs. (1b) and (12) into eqs. (2b) and

$$\mu = 8\pi an \left[1 + 4 \left(\frac{8}{3\sqrt{\pi}} + c_{\text{ip}} \right) \sqrt{a^3 n} \right], \quad (29)$$

respectively. Finally using the thermodynamic relation $\mu = \partial E / \partial N$ to integrate eq. (29) over N , we arrive at eq. (2a). Equation (2) with eq. (26) are correct up to the next-to-the-leading order.

Finally, let us estimate c_{ip} by using two different approximations. First, retaining only the first two terms on the right-hand side of eq. (23) yields a cubic equation for $y = y(x)$, which is defined in eq. (25), as

$$y^3 - 2by^2 + (b^2 + 2)y - 2(b + 5) = 0. \quad (30)$$

It has a single real solution $y = y(x)$ for $0 \leq x \leq \infty$ that approaches 0 continuously as $x \rightarrow \infty$, in accordance with $\Delta_{\vec{p}}^{(\text{ip})}(i\varepsilon_\ell) \rightarrow 0$ as $|\varepsilon_\ell| \rightarrow \infty$. Substituting it into eq. (26) and performing the integration, we obtain $c_{\text{ip}} = 0.412$. Hence, the factor $128/15\sqrt{\pi} = 4.81$ in eq. (1a) is changed into $(16/5)(8/3\sqrt{\pi} + c_{\text{ip}}) = 6.13$ in eq. (2a). Second, eq. (13b) may be generalized to the n th-order *improper* terms ($n = 3, 4, \dots$) originating from the particle-hole and particle-particle bubble diagrams as

$$\Sigma_{\vec{p}}^{(n\text{ip})} = \Delta_{\vec{p}}^{(n\text{ip})} = \left(1 + \frac{1}{2^{n-1}} \right) (2Un_0)^n f_{\vec{p}}^{n-1}, \quad (31)$$

where $f_{\vec{p}}$ is defined by eq. (14). Collecting the series for $n = 3, 4, \dots, \infty$ together with eq. (13a) of the second order, we obtain

$$\Delta_{\vec{p}}^{(\text{ip})} = \frac{(2Un_0)^2 f_{\vec{p}}}{1 - 2Un_0 f_{\vec{p}}} + \frac{2(Un_0)^2 f_{\vec{p}}}{1 - Un_0 f_{\vec{p}}} - 4(Un_0)^2 f_{\vec{p}} \\ = -\frac{(2Un_0)^2}{\eta_{\vec{p}} + 2Un_0 + \Delta_{\vec{p}}^{(\text{ip})}} - \frac{2(Un_0)^2}{\eta_{\vec{p}} + Un_0 + \Delta_{\vec{p}}^{(\text{ip})}} \\ + \frac{4(Un_0)^2}{\eta_{\vec{p}} + \Delta_{\vec{p}}^{(\text{ip})}}. \quad (32)$$

This self-consistent equation for $\Delta_{\vec{p}}^{(\text{ip})}$ is transformed in terms of the quantities in eq. (25) into

$$y^4 - 3(b-1)y^3 + (3b^2 - 6b + 4)y^2 \\ - (b^3 - 3b^2 + 6b + 4)y + 2(b^2 + 2b - 4) = 0, \quad (33)$$

which also has a single real solution $y = y(x)$ for $0 \leq x \leq \infty$ that approaches 0 continuously as $x \rightarrow \infty$. Substituting it into eq. (26), we obtain $c_{\text{ip}} = 0.563$. Thus, we may expect that c_{ip} is of order 1. The exact value of c_{ip} can only be reached by collecting all the leading-order *improper* terms, however, meaning that further studies are required for a quantitative estimate of c_{ip} . On the other hand, a diffusion Monte Carlo study was performed on eq. (1);¹³ the data in Table I of ref. 13 for $a^3 n \leq 10^{-2}$

yield the value 4.3(1), rather than $128/15\sqrt{\pi} = 4.81$ from Eq. (1a), thus showing a clear deviation from the Lee-Huang-Yang expression beyond the numerical uncertainty. On the other hand, an experiment on eq. (1a) in a trap potential reported a value 4.5(7),¹⁴ which still requires improvement for the present purpose, however.

Acknowledgments

We are grateful to D. Hirashima for a useful discussion on the diffusion Monte Carlo study. This work is supported in part by JSPS Grant (C) No. 22540356.

-
- ¹ T. D. Lee, K. Huang, and C. N. Yang: Phys. Rev. **106** (1957) 1135.
 - ² N. N. Bogoliubov: J. Phys. (USSR) **11** (1947) 23.
 - ³ A. A. Abrikosov, L. P. Gorkov, and I. E. Dzyaloshinski: *Methods of Quantum Field Theory in Statistical Physics* (Prentice Hall, Englewood Cliffs, N.J., 1963).
 - ⁴ A. L. Fetter and J. D. Walecka: *Quantum Theory of Many-Particle Systems* (McGraw-Hill, New York, 1971).
 - ⁵ M. Girardeau and R. Arnowitt: Phys. Rev. **113** (1959) 755.
 - ⁶ A. Griffin: Phys. Rev. B **53** (1996) 9341.
 - ⁷ N. M. Hugenholtz and D. Pines: Phys. Rev. **116** (1959) 489.
 - ⁸ P. C. Hohenberg and P. C. Martin: Ann. Phys. (N.Y.) **34** (1965) 291.
 - ⁹ T. Kita: Phys. Rev. B **80** (2009) 214502.
 - ¹⁰ T. Kita: Phys. Rev. B **81** (2010) 214513.
 - ¹¹ T. Kita: J. Phys. Soc. Jpn. **80** (2011) 084606.
 - ¹² J. M. Luttinger and J. C. Ward: Phys. Rev. **118** (1960) 1417.
 - ¹³ S. Giorgini, J. Boronat, and J. Casulleras: Phys. Rev. A **60** (1999) 5129.
 - ¹⁴ N. Navon, S. Piatecki, K. Günter, B. Rem, T. C. Nguyen, F. Chevy, W. Krauth, and C. Salomon: Phys. Rev. Lett. **107** (2011) 135301.